




Revista Latinoamericana de Psicología

<http://revistalatinoamericanadepsicologia.konradlorenz.edu.co/>



ORIGINAL

Depression and associated psychosocial factors following a legal abortion in Mexico

Luciana Ramos-Lira^{a,*}, Karla Flores Celis^b, Midiam Moreno López^a,
María Teresa Saltijeral^a, Fernando Wagner^c

^a Department of Epidemiologic and Psychosocial Research, National Institute of Psychiatry Ramón de la Fuente Muñiz, Mexico

^b Facultad de Medicina, Universidad Nacional Autónoma de México, México

^c School of Social Work, University of Maryland, Maryland, USA

Received 22 June 2022; accepted 17 January 2023

KEYWORDS

Induced legal abortion, depression, psychosocial factors, contextual factors, abortion stigma, Latin America

Abstract Introduction: Scientific evidence from the United States and European countries shows that women who have had an induced abortion are not more likely to become depressed and several factors may confound this outcome. In contrast, in the case of Latin America, there are practically no studies in this regard due to restrictive legislation prevailing in the region. This paper aims to determine the prevalence of a probable major depressive episode (PMDE) in women who have legally terminated a pregnancy by way of public service in Mexico City and whether there are any psychosocial factors reported by the international literature associated with this outcome. **Method:** In a cross-sectional study, 274 women aged 15 years or older were interviewed two weeks after undergoing a medical abortion between November 2018 and November 2019. The Center for Epidemiological Studies Depression Scale 35-item version (CES-D-R35) was used to measure the PMDE in a structured interview. **Results:** The prevalence of a PMDE was 15.8%. Multivariate logistic regression was used for adjusted analysis. Perceived abortion stigma ($OR = 6.74$, $95\% CI = 3.29-13.82$), child sexual abuse ($OR = 2.23$, $95\% CI = 1.01-4.93$), and previous childless pregnancies ($OR = 6.07$, $95\% CI = 1.52-24.21$) were associated with PMDE. **Conclusions:** The prevalence of PMDE is similar to or lower than that reported in studies with women who continued a pregnancy; post-abortion counseling and clinical considerations should include the impact that stigma and gender-based violence have on women's mental health.

© 2023 Fundación Universitaria Konrad Lorenz. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Depresión y factores psicosociales asociados tras un aborto legal en México

PALABRAS CLAVE

Aborto legal inducido, depresión, factores psicosociales, factores contextuales, estigma por abortar, Latinoamérica

Resumen Introducción: Evidencia científica proveniente de los Estados Unidos y los países europeos indica que las mujeres que han tenido un aborto inducido no tienen más probabilidades de deprimirse y que hay varios factores que pueden intervenir para confundir este resultado. En el caso de América Latina, prácticamente no existen estudios al respecto debido a las legislaciones restrictivas imperantes. El objetivo de este artículo es determinar la prevalencia de un probable episodio depresivo mayor (PEDM) en mujeres que interrumpieron legalmente un embarazo en un servicio público en la Ciudad de México,

* Corresponding author.

E-mail: luciana.ramos.lira@gmail.com

<https://doi.org/10.14349/rlp.2023.v55.8>

0120-0534/© 2023 Fundación Universitaria Konrad Lorenz. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

y si hay algunos factores psicosociales reportados por la literatura internacional asociados con el resultado. **Método:** En un estudio transversal, se entrevistó a 274 mujeres de 15 años o más dos semanas después de someterse a un aborto médico entre noviembre del 2018 y noviembre del 2019. Se utilizó la versión de 35 reactivos de la Escala de Depresión del Centro para Estudios Epidemiológicos (CES-D-R35) para medir el PEDM en una entrevista estructurada. **Resultados:** La prevalencia de PEDM fue de 15,8%. Se utilizó una regresión logística multivariada para el análisis ajustado. El estigma percibido por el aborto ($OR = 6.74$, $IC\ 95\% = 3.29-13.82$), el abuso sexual infantil ($OR = 2.23$, $IC\ 95\% = 1.01-4.93$) y los embarazos previos sin hijos ($OR = 6.07$, $IC\ 95\% = 1.52-24.21$) se asociaron con un PMDE. **Conclusiones:** La prevalencia de PEDM es similar o menor que la reportada en estudios con mujeres que continuaron un embarazo; el asesoramiento posaborto y las consideraciones clínicas deben incluir el impacto que el estigma y la violencia de género tienen en la salud mental de las mujeres.

© 2023 Fundación Universitaria Konrad Lorenz. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Most research on the effects of induced abortion on women's mental health has been conducted in high-income countries (APA, 2008; Biggs et al., 2017; National Collaborating Centre for Mental Health, 2011; van Ditzhuijzen, 2019). However, rigorous evidence shows that abortion is not causally associated with adverse mental health outcomes (Biggs et al., 2018; Lewis & Dave, 2021; Major et al., 2009; Steinberg et al., 2014; Stotland, 2019; van Ditzhuijzen et al., 2017).

Depression has been one of the most researched outcomes subsequent to an abortion. Notwithstanding, the evidence has revealed that several factors could be confounding this association (Boersma et al., 2014; Charles et al., 2008; Faure & Loxton, 2003; Foster et al., 2015; Pederesen, 2008; Rees & Sabia, 2007). For example, a history of gender-based violence (Major et al., 2009; Roberts et al., 2014; Steinberg et al., 2016; Oram et al., 2017; Zia et al., 2021), the previous presence of a mental health problem (Steinberg & Rubin, 2014; van Ditzhuijzen et al., 2017), or the perceived stigma of abortion (Major et al., 2009; López et al., 2019; Ramos-Lira et al., 2021; Sorhaindo et al., 2014; Steinberg et al., 2016; Zia et al., 2021).

In light of this background, we hypothesise that women who abort may have mental health conditions that are not the result of an abortion procedure and that other prior or concomitant factors may lead to these mental health outcomes. Likewise, it is relevant to carry out studies on the subject in Latin America, taking into account that although this region has highly restrictive abortion laws, it also has the highest rates of induced abortion worldwide (Galli, 2020; Singh et al., 2018). Consequently, there are many barriers to monitoring and researching this issue (Biggs et al., 2020; Boersma et al., 2014; Cedeño & Tena Guerrero, 2022; Maroto-Vargas, 2010).

In 2021, the Mexican Supreme Court of Justice declared it unconstitutional to criminalize abortion in an absolute way. Legal termination of pregnancy by public services exists in ten states and is available up to the 12th week of gestation. Moreover, services have existed in Mexico City since 2007, when abortion was decriminalised, and are currently the most consolidated in the country.

This paper aims to determine the prevalence of probable major depressive episode (PMDE) in women who legally terminated a pregnancy by way of public service in Mexico City. It also seeks to identify whether there are any associated psychosocial and contextual factors.

Method

Type of study and participants

The research was cross-sectional and ex-post-facto, involving 274 Mexican women who returned for a follow-up medical examination two weeks after a medical abortion provided by a public service of Legal Termination of Pregnancy (LTP) in Mexico City. Eligibility for the study included being 15 years of age or older and agreeing to participate in an interview. Exclusion criteria included women a) that had performed the procedure with manual vacuum aspiration (MVA), b) that had a severe intellectual or motor disability that would not allow them to answer the questionnaire, or c) that had attended the service for some other legal reason.

Measures

A structured interview included questions regarding age, education, occupation, socioeconomic status, number of previous pregnancies, previous abortions, number of children, and the presence of a partner.

Depression was measured using the Revised Version of the Depression Scale of the Center for Epidemiological Studies (CES-D-R; González-Forteza et al., 2008). The instrument includes 35 questions with five closed answer options referring to the number of days in which symptoms persisted (0, 1-2, 3-4, 5-7, 8-14), with regard to the last 14 days. It was created in 1977 (Radloff, 1977) and revised in 2004 (Eaton et al., 2004) for measuring the symptoms defined by the Diagnostic and Statistical Manual of Mental Disorders, 4th Edition (DSM-IV; APA, 1994) of a major depressive episode. In the present study, the instrument obtained an $\alpha = 0.95$. Women were classified using an algorithm for a probable major depressive episode (PMDE) that evaluates the presence or absence of symptoms in each of the nine criteria for a major depressive episode. Hence, a score threshold is not used for depression screening in our study.

To assess whether participants had experienced previous depression, they were asked the following question after completing the CES-D-R scale: "In the last 12 months or from (the current month but referring to the last year) and




Figure 1. Sample recruitment description

until you attended the LTP service, have you had symptoms like the ones I just mentioned at the same time and with such intensity and duration that you would say you were depressed?” Again answer options were dichotomous: yes or no.

Child Sexual Abuse was assessed with one question: “Before the age of 15, has someone, whether a family member or anyone else, ever stroked, touched, or caressed some part of your body; or did they make you touch any part of their body, or did they have sex with you when you were very young or against your will?” Answer options were, again, dichotomous: yes or no. Another question included was regarding forced sex: “Starting at age 15, did anyone force you to have sex or when you did not want to?” Once again, the answer options were a simple yes or no (Ramos-Lira et al., 2001).

Intimate Partner Violence (IPV) during the last year prior to going to the LTP service was evaluated with four items, taking as a reference the current or the latest partner (Lira & Méndez, 2008). Two items assessed psychological abuse (You were yelled at or insulted; threatened to be physically harmed, for example, to be hurt or even killed), a second item considered physical abuse (You were physically abused; this includes being slapped, kicked or hit). Another one referred to sexual abuse (He has had sex with you by force, even though you didn’t want to), and finally, there was an economic violence item (He has complained about how you spend money). As mentioned above, the response options were all dichotomous. For the analyses, a single IPV variable was created considering the presence of at least one of the four items ($\alpha = 0.72$); all answers were dichotomous.

The Individual Abortion Stigma Scale (ILAS; Cockrill et al., 2013) comprises 20 items and measures four dimen-

sions: Worries about judgment, Isolation, Self-judgment, and Community condemnation. These items have four Likert-type response options for the first two dimensions (from 1 = *nothing* to 4 = *a lot*, subscale range 1-4), while the third and fourth have five options (from 1 = *strongly disagree*, to 5 = *strongly disagree*; and from 1 = *none*, to 5 = *most people*, respectively, subscale range 1-5). Subscale scores were obtained by adding each item and dividing the sum by the number of questions. The range for the total score was 1-4.35. There was no set cut-off point. The scale showed good reliability ($\alpha = 0.88$ to 0.98).

At the end of the interview, if a participant had presented a history of depression, current symptoms, a severe case of violence within the last 12 months, or if she had requested care, she was offered a referral to the hospital’s psychiatry or psychology service or to another institution.

Procedure

Interviews with users of the LTP service took place between November 2018 and November 2019. Five interviewers participated in data collection; all were psychologists and had previously participated in specialised training.

A researcher and the social worker assigned to the corresponding public service invited all women attending follow-up consultations to participate. Participants signed an informed consent letter and were invited to a quiet, private location to be interviewed directly. The questionnaire was then applied during the waiting time for the follow-up ultrasound.

During the selected time for data collection, 714 women attended the service. Of these, four requested an interruption for reasons associated with a danger to their health; 14 voluntarily withdrew from the service and did not complete

the process; the gestation time of 35 women was more than 12 weeks advanced at the time of ultrasound, and they could not receive the termination, and 93 others performed the procedure with MVA. Of the 568 who performed the procedure with medication, 460 were eligible with regard to the time when the research team showed up at the service to conduct the interviews. Of those eligible, only 312 returned for the discharge appointment and they were invited to participate; of these, 274 accepted the interview (87.8%) (Figure 1).

Ethical considerations

This research project was evaluated and approved by the Ethics Committee of the Instituto Nacional de Psiquiatría Ramón de la Fuente Muñiz, reference CEI/C/001/2015, and by the Research and Bioethics Subcommittee of the Secretaría de Salud of Mexico City (SSDF/ DGPCS/DEI/SECI/ JUDI /1279/18).

Statistical analyses

Analyses were performed in STATA 14.0. The descriptive data of the variables were analysed in absolute frequencies. Bivariate analyses were performed between psychosocial variables and PMDE using Student's χ^2 or T-tests with a $p < .05$. Odds ratios (ORs) and confidence intervals (95% CI) were estimated using a logistic regression model, including variables significant in the bivariate analysis and adjusted for age, education, occupation, socioeconomic status, and PMDE as the outcome.

Results

Most women were between 20 and 29 years old (57.7%). Half had completed secondary school (50.4%), and approximately the same proportion reported having a paid activity. Most participants had a very low or low socioeconomic status (89.4%). Almost four out of five mentioned having a partner (78.1%); half reported cohabiting with him.

65.6% had been pregnant before, and 10.3% had undergone a previous induced or spontaneous abortion. In addition, 32.1% reported child sexual abuse (before age 15), and 12.1% were forced to have sex after that age. Two out of five (39.9%) reported depression in the year prior to the abortion; 47.1% had experienced IPV in the past 12 months, with emotional violence being the most common form. 15.8% of the sample presented a PMDE. Perceived abortion stigma was not very high, as observed in the scale's total score, with worries concerning judgment and community condemnation being the highest dimensions (Table 1).

Regarding reproductive history, Table 2 shows that women with PMDE were more likely to have been pregnant before without having had children (due to induced abortion/miscarriage) than those without PMDE (20.9% vs. 8.3%). Also, those with PMDE were more likely to report child sexual abuse compared to those without PMDE (53.5% vs. 28.1%) ($\chi^2(1) = 10.72, p = 0.001$). Women with PMDE were also more

likely to have experienced forced sex after age 15 (23.3% vs. 10%) ($\chi^2(1) = 5.93, p = 0.015$) as well as IPV within the past 12 months (65.1% vs. 43.7%) ($\chi^2(1) = 6.68, p = 0.010$). Moreover, those with PMDE were more likely to report depression during the 12 months before pregnancy (58.1% vs. 36.4%) ($\chi^2(1) = 7.13, p = 0.008$).

Women with PMDE reported higher averages of abortion stigma in the Total Score compared to those without PMDE (2.99 vs. 2.29), ($t(270) = 6.86, p < .001$), and likewise in all four dimensions compared to those without PMDE: Worries about judgment (2.67 vs. 2.06), ($t(270) = 4.48, p < .001$); Isolation (2.34 vs. 1.84), ($t(270) = 3.71, p < .001$); Self-judgment (3.47 vs. 2.33), ($t(270) = 6.96, p < .001$); and Community condemnation (3.49 vs. 2.92), ($t(270) = 2.58, p = 0.010$).

As shown on Table 3, unadjusted estimates of reproductive history, child sexual abuse, forced sex after the age of 15, any IPV, self-report of depression, and abortion stigma were associated with PMDE. Subsequently, in the final logistic regression model and adjusting for sociodemographic characteristics, higher ORs were found among women who had been pregnant but had not borne children ($OR = 6.07$; 95% $CI = 1.52-24.21$); likewise for those who reported child sexual abuse ($OR = 2.23$; $CI 95\% = 1.01-4.93$) and for those with higher levels of perceived abortion stigma ($OR = 6.74$; 95% $CI = 3.29-13.82$).

Discussion

The findings contribute to understanding the abortion situation in Latin America, where there are practically no studies on the subject due to the legal restrictions associated with its practice. The prevalence of PDME was 15.8%, similar to or lower than that reported in meta-analyses of studies of women who continued a pregnancy, which ranges from 15.0% to 20.7% (Dadi et al., 2020; Yin et al., 2021).

Abortion-related stigma was the most crucial factor in the PMDE model. This result is consistent with results reported in other international studies (Biggs et al., 2020; O'Donnell et al., 2018; Steinberg et al., 2016) and with the proposals of authors such as Rostagnol (2014), who points out that while the decriminalisation of abortion solves the problem of unsafe abortions, it does not generate automatic changes in moral beliefs.

Although attitudes toward abortion tend to be more favourable after legalisation, and the circumstances for performing it are more likely to be considered reasonable if the law allows it (Sorhaindo et al., 2014), abortion stigma has lasting effects on people's lives (Assis & Erdman, 2021). This stigma is a central component of anti-choice narratives, in which abortion is constructed as a shameful, immoral, and deviant practice (Amuchástegui et al., 2015; Cullen & Korolczuk, 2019; Zia et al., 2021).

Abortion stigma can produce feelings of guilt and shame that often lead women to keep the experience secret. (Cedeño & Tena Guerrero, 2022; Hanschmidt et al., 2016; O'Donnell et al., 2018; Shellenberg et al., 2011; Sorhaindo et al., 2014; Zia et al., 2021). It also often causes fear of being judged by relatives and significant others, which can ultimately lead to psychological distress (Biggs et al., 2020; Hanschmidt et al., 2016) or even symptoms of depression (Steinberg et al., 2016). This concern about

Table 1. Descriptive statistics of the study sample (n = 274)

Variables		n	%
Sociodemographic characteristics			
Age (years)	15-19	55	20.1
	20-24	91	33.2
	25-29	67	24.5
	30-34	37	13.5
	35 or more	24	8.8
Educational attainment	Undergraduate or graduate	60	21.9
	High school	138	50.4
	Middle school	66	24.1
	Primary	10	3.6
Occupation	Remunerated activity	137	50.0
	Unpaid activity	81	29.6
	Student	56	20.4
Socioeconomic status*	Middle	29	10.6
	Low	152	55.5
	Very low	93	33.9
Currently has an intimate partner	Yes	214	78.1
Psychosocial factors			
Reproductive history	No previous pregnancies	94	34.4
	Previous pregnancies without children **	28	10.3
	Previous pregnancies with children	151	55.3
Experiences of sexual violence	Any child sexual abuse (before the age of 15)	87	32.1
	Any forced sexual intercourse (from the age of 15)	33	12.1
Self-reported depression within the 12 months prior to pregnancy	Yes	108	39.9
Intimate partner violence within the last 12 months**	Emotional	98	36.0
	Sexual/physical	55	20.2
	Economic	73	26.8
	Any type of intimate partner violence	128	47.1
PMDE	No	229	84.2
	Yes	43	15.8
		n	Mean
Perceived abortion stigma	Worries about judgment	274	2.15
	Isolation	274	1.92
	Self-judgment	274	2.51
	Community condemnation	274	3.01
	Total score	274	2.40

* Evaluated by a social worker with the classification of the LTP service, **Add up to more than one hundred per cent because categories are not mutually exclusive.

being judged also interferes with the search for abortion services (Hernández-Rosete & Estrada-Hipólito, 2019; Turan & Budhwani, 2021).

Having a history of abortions and the absence of children was a risk factor for depression that may be related to idealised beliefs about motherhood and stigmatised precepts about abortion. Women who have decided to terminate several pregnancies and do not have children may feel selfish or immoral because they perceive that they are challenging family expectations and cultural norms regarding motherhood and womanhood (Kumar et al., 2009; Orihuela-Cortés et al., 2022; Russo, 1976). Stigmatisation provokes a narrative in which women consider it valid to have a single abortion for a justified reason (Amuchástegui et al., 2015). Women with more than one induced abortion are regarded as careless or promiscuous.

In addition to all of the above, this study evidences the relevance of child sexual abuse. The literature regarding this issue shows that this is a common risk factor for the development of mental health disorders in general (Hailes et al., 2019) and depression in particular (Li et al., 2020), as well as for subsequent sexual victimisation, unwanted pregnancy (Sumner et al., 2015), reduction of contraceptive use (Grose et al., 2021), and repetition of induced abortion (Fisher et al., 2005).

Limitations and suggestions

Despite the strengths of this study, it is important to note its limitations:

1. The design was cross-sectional, so the causal nature of the relationships examined cannot be determined. A prospective study could provide better information to discriminate the magnitude of the different factors considered in the presence of PMDE.
2. Only one service participated in the study, which does not necessarily mean that this population is representative of other LTP clinics.
3. Interviews were conducted exclusively with women who returned to their discharge appointment following a medical abortion because this is the most frequent procedure afforded by LTP services in Mexico City (78.95% vs. 21.04% for MVA). Unfortunately, we do not know how patients who returned for their follow-up appointments differ from those who did not return.
4. There was no comparison group, and several variables such as violent experiences and self-reported depression included dichotomous answers to measure complex phenomena.

Table 2. Bivariate analysis of psychosocial variables and PMDE ($n = 274$)

Variables	PMDE			
	No		Yes	
	<i>n</i>	%	<i>n</i>	%
Reproductive history				
No previous pregnancies	84	36.8	9	20.9
Previous pregnancies without children **	19	8.3	9	20.9
Previous pregnancies with children	125	54.8	25	58.1
Currently has an intimate partner	182	79.5	31	72.1
Experiences of sexual violence				
Any child sexual abuse *	64	28.1	23	53.5
Any forced sexual intercourse**	23	10.0	10	23.3
Self-reported depression within the 12 months prior to pregnancy*	83	36.4	25	58.1
Intimate partner violence within the last 12 months				
Emotional**	76	33.2	22	51.2
Sexual/physical*	38	16.6	17	39.5
Economic	59	25.8	14	32.6
Any intimate partner violence *	100	43.7	28	65.1
Perceived abortion stigma	Mean (SD)		Mean (SD)	
Worries about judgment***	2.06 (.82)		2.67 (.87)	
Isolation***	1.84 (.78)		2.34 (.94)	
Self-judgment***	2.33 (1.0)		3.47 (.82)	
Community condemnation***	2.92 (1.3)		3.49 (1.4)	
Total score***	2.29 (.62)		2.99 (.61)	

* $\chi^2 p < 0.01$, ** $\chi^2 p < 0.05$, *** Student's $T p < 0.01$

Table 3. Logistic regression models of factors associated with PMDE (*n* = 269)

Variables in the model	PMDE					
	OR Crude	<i>p</i>	95% CI	Adjusted OR*	<i>p</i>	95% CI
Age (continuous)	0.97	0.312	0.92-1.03	0.94	0.141	0.86-1.02
Educational attainment						
<i>High school or more</i>	1.00	(reference)		1.00	(reference)	
<i>Middle school or less</i>	1.51	0.245	0.75-3.02	1.22	0.667	0.50-2.97
Occupation						
Remunerated activity	1.00	(reference)		1.00	(reference)	
<i>Unpaid activity</i>	1.65	0.183	0.79-3.46	1.74	0.242	0.69-4.42
<i>Student</i>	1.29	0.562	0.54-3.09	1.10	0.880	0.32-3.84
Socioeconomic status						
<i>Middle</i>	1.00	(reference)		1.00	(reference)	
<i>Low</i>	1.65	0.438	0.46-5.90	1.19	0.819	0.27-5.25
<i>Very low</i>	1.60	0.483	0.43-5.99	1.25	0.782	0.26-5.96
Reproductive history						
<i>No previous pregnancies</i>	1.00	(reference)		1.00	(reference)	
<i>Previous pregnancies without children</i>	4.42	0.006	1.55-12.63	6.07	0.011	1.52-24.21
<i>Previous pregnancies with children</i>	1.87	0.131	0.83-4.20	1.77	0.341	0.55-5.77
Currently has an intimate partner (vs. no partner)	0.67	0.283	0.32-1.40	1.07	0.885	0.41-2.81
Childhood sexual abuse (vs. no)	2.95	0.001	1.52-5.73	2.23	0.048	1.01-4.93
Forced sexual relation (vs. no)	2.71	0.018	1.19-6.21	2.06	0.157	0.76-5.59
Any intimate partner violence (vs. no)	2.41	0.011	1.22-4.75	1.07	0.883	0.45-2.53
Self-report of depression (vs. no)	2.43	0.009	1.25-4.71	1.55	0.291	0.69-3.49
Total score perceived abortion stigma (continuous)	5.88	< 0.001	3.19-10.81	6.74	< 0.001	3.29-13.82

* Multiple logistic regression analysis with method Enter, all the variables indicated on the Table were introduced. The model was significant and explained 30% of the variance.

5. The report of a low and very low socioeconomic level (89%), considering the high educational level of the participants, is striking. In this sense, it is crucial to consider a possible bias in this variable given that the social worker of the service assigned the participants' socioeconomic level based on an institutional criterion that allows for the interruption of pregnancy to be free.

It is urgent to promote a "social" decriminalisation of abortion, that is, to accept that induced abortion is a frequent procedure and, most importantly, a right for women who wish to terminate a pregnancy for whatever reason. Moreover, making women feel less stigmatised, decreasing misinformation related to the termination of a pregnancy, improving their perception of self-efficacy and self-care as well as detecting experiences of violence could reduce the emotional burden related to making decisions concerning their sexual and reproductive health.

These findings coincide with the arguments on reproductive autonomy put forth by the Mexican Supreme Court of Justice in 2021. Obstacles in women's sexual and reproductive trajectory for making free decisions disrupt their care structures, where preventing an unintended pregnancy becomes very unlikely or controllable for them. Therefore, abortion can be seen as a therapeutic intervention that guarantees their right to health and the free development of their personality.

Services must be available, accessible, and of high technical quality. They should include post-abortion counseling and clinical considerations with regard to the impact of stigma and violence on women's mental health and choices. Establishing intra-hospital pathways between different services that guarantee integral health when someone requires this type of care is crucial. Sensitive personnel with active listening skills are also required to establish a doctor-patient relationship focused on the needs of the women in each case. Finally, it is particularly relevant to analyse whether a brief clinical intervention to explore stigmatising ideas could help to anticipate depressive symptoms following the termination of a pregnancy.

Conflict of interest

There is no conflict of interest.

Funding

This work was supported by the Consejo Nacional de Ciencia y Tecnología (CONACYT) under Grant CB-2015-01-0254877.

Acknowledgments

To the scholarship holders of the Psychology degree programme who carried out the fieldwork: Haidee Bautista Reyes, Ambar Rodríguez Flores, and Rocío Rojas Hernández. To Catalina González-Forteza, Raffaella Schiavon, and María Elena Collado for their suggestions and comments. To Olivia Ortiz for collaborating with the research team and the participants who required it. To doctors Juan Carlos de la Cerda Ángeles and Emilio Exaire for their support in carrying out this study in the LTP service. To nurse Angélica Álvarez González, to psychologist Magdalena Hernández Espinoza, and very significantly to social worker Rocío Sánchez Hurtado for her valuable synergy. Karla Flores Celis thanks the National Council of Science of Technology (CONACyT) for the doctoral scholarship awarded (CVU 624806).

References

- American Psychiatric Association (1994). *Diagnostic and Statistical Manual of Mental Disorders* (4th ed.). [DSM-IV]. American Psychiatric Publishing, Inc.
- American Psychological Association (2008). *Report of the task force on mental health and abortion*. American Psychological Association. <https://www.apa.org/pi/women/programs/abortion/mental-health.pdf>
- Amuchástegui, A., Flores, Edith., & Aldaz, E. (2015). Disputa social y disputa subjetiva. Religión, género y discursos sociales en la legalización del aborto en México. *Revista de Estudios de Género La Ventana*, 5(41), 153-195. <https://doi.org/10.32870/lv.v5i41.4315>
- Assis, M. P., & Erdman, J. N. (2021). Abortion rights beyond the medico-legal paradigm. *Global Public Health*, 17(10), 2235-2250. <https://doi.org/10.1080/17441692.2021.1971278>
- Biggs, M. A., Brown, K., & Foster, D. G. (2020). Perceived abortion stigma and psychological well-being over five years after receiving or being denied an abortion. *Plos One*, 15(1), e0226417. <https://doi.org/10.1371/journal.pone.0226417>
- Biggs, M. A., Gould, H., Barar, R. E., & Foster, D. G. (2018). Five-Year Suicidal Ideation Trajectories Among Women Receiving or Being Denied an Abortion. *The American Journal of Psychiatry*, 175(9), 845-852. <https://doi.org/10.1176/appi.ajp.2018.18010091>
- Biggs, M. A., Upadhyay, U. D., McCulloch, C. E., & Foster, D. G. (2017). 'Women's mental health and well-being 5 years after receiving or being denied an abortion: A prospective, longitudinal cohort study. *JAMA Psychiatry*, 74(2), 169-178. <https://doi.org/10.1001/jamapsychiatry.2016.3478>
- Boersma, A. A., Van Den Berg, D., Van Lunsen, R. H., & Laan, E. T. (2014). Induced abortion is not associated with a higher likelihood of depression in Curaçao women. *The European Journal of Contraception & Reproductive Health Care*, 19(5), 359-367. <https://doi.org/10.3109/13625187.2014.927422>
- Cedeño, L., & Tena Guerrero, O. (2022). Estigma y empoderamiento posterior al aborto en mujeres mexicanas. *Iberoforum. Revista de Ciencias Sociales*, 2(1), 1-46. <https://doi.org/10.48102/if.2022.v2.n1.212>
- Charles, V. E., Polis, C. B., Sridhara, S. K., & Blum, R. W. (2008). Abortion and long-term mental health outcomes: A systematic review of the evidence. *Contraception*, 78(6), 436-450. <https://doi.org/10.1016/j.contraception.2008.07.005>
- Cockrill, K., Upadhyay, U. D., Turan, J., & Greene-Foster, D. (2013). The stigma of having an abortion: Development of a scale and characteristics of women experiencing abortion stigma. *Perspectives on Sexual and Reproductive Health*, 45(2), 79-88. <https://doi.org/10.1363/4507913>
- Cullen, P., & Korolczuk, E. (2019). Challenging abortion stigma: Framing abortion in Ireland and Poland. *Sexual and Reproductive Health Matters*, 27(3), 6-19. <https://doi.org/10.1080/26410397.2019.1686197>
- Dadi, A. F., Miller, E. R., Bisetegn, T. A., & Mwanri, L. (2020). Global burden of antenatal depression and its association with adverse birth outcomes: An umbrella review. *BMC Public Health*, 20(1), 1-16. <https://doi.org/10.1186/s12889-020-8293-9>
- Eaton, W.W., Smith, C., Ybarra, M., Muntaner, C., & Tien, A. (2004). Center for Epidemiologic Studies Depression Scale: Review and Revision (CESD and CESD-R). In Maruish M. E. (Ed.), *The use of psychological testing for treatment planning and outcomes assessment: Instruments for adults* (pp. 363-377). Routledge, NY. <https://doi.org/10.4324/9781410610614>
- Faure, S., & Loxton, H. (2003). Anxiety, depression and self-efficacy levels of women undergoing first trimester abortion. *South African Journal of Psychology*, 33(1), 28-38. <https://doi.org/10.1177/0081246303300104>
- Fisher, W. A., Singh, S. S., Shuper, P.A., Carey, M., Otchet, F., MacLean-Brine, D., Dal Bello, D., & Gunter, J. (2005). Characteristics of women undergoing repeat induced abortion. *Canadian Medical Association Journal*, 172(5), 637-641. <https://doi.org/10.1503/cmaj.1040341>
- Foster, D. G., Steinberg, J. R., Roberts, S.C., Neuhaus, J., & Biggs, M. A. (2015). A comparison of depression and anxiety symptom trajectories between women who had an abortion and women denied one. *Psychological Medicine*, 45(10), 2073-2082. <https://doi.org/10.1017/S0033291714003213>
- Galli, B. (2020). Challenges and opportunities for access to legal and safe abortion in Latin America based on the scenarios in Brazil, Argentina, and Uruguay. *Cadernos De Saúde Pública*, 36(1), e00168419. <https://doi.org/10.1590/0102-311X00168419>
- González-Forteza, C., Jiménez-Tapia, J. A., Ramos-Lira, L., & Wagner, F. A. (2008). Aplicación de la Escala de Depresión del Center of Epidemiological Studies en adolescentes de la ciudad de México. *Salud Pública de México*, 50(4), 292-299. <https://doi.org/10.1590/S0036-36342008000400007>
- Grose, R. G., Chen, J. S., Roof, K. A., Rachel, S., & Yount, K. M. (2021). Sexual and reproductive health outcomes of violence against women and girls in lower-income countries: A review of reviews. *The Journal of Sex Research*, 58(1), 1-20. <https://doi.org/10.1080/00224499.2019.1707466>
- Hailes, H. P, Yu, R., Danese, A., & Fazel, S. (2019). Long-term outcomes of childhood sexual abuse: An umbrella review. *Lancet Psychiatry*, 6(10), 830-839. [https://doi.org/10.1016/S2215-0366\(19\)30286-X](https://doi.org/10.1016/S2215-0366(19)30286-X)
- Hanschmidt, F., Linde, K., Hilbert, A., Riedel-Heller, S. G., & Kersting, A. (2016). Abortion stigma: A systematic review. *Perspectives on Sexual and Reproductive Health*, 48(4), 169-177. <https://doi.org/10.1363/48e8516>
- Hernández-Rosete, D., & Estrada-Hipólito, R. (2019). Dificultades de acceso al aborto en contextos de interrupción legal del embarazo: narrativa de estudiantes de bachillerato en una comunidad campesina de México. *Cadernos De Saúde Pública*, 35(3), 1-11. <https://doi.org/10.1590/0102-311X00046218>
- Kumar, A., Hessini, L., & Mitchell, E. M. (2009). Conceptualising abortion stigma. *Culture, Health & Sexuality*, 11(6), 625-639. <https://doi.org/10.1080/13691050902842741>
- Lewis, E. G., & Dave, U. (2021). Long-term mental health & financial outcomes associated with abortion access and denial: A systematic review. *Journal of Clinical Obstetrics & Gynecology*, 31(3), 97-99. <https://doi.org/10.5336/jcog.2021-85039>
- Li, D., Chu, C. M., & Lai, V. (2020). A developmental perspective on the relationship between child sexual abuse and depression:

- A systematic and meta-analytic review. *Child Abuse Review*, 29(1), 27-47. <https://doi.org/10.1002/car.2592>
- Lira, L. R., & Méndez, M. T. S. (2008). ¿Violencia episódica o terrorismo íntimo? Una propuesta exploratoria para clasificar la violencia contra la mujer en las relaciones de pareja. *Salud Mental*, 31, 469-478. http://www.revistasaludmental.mx/index.php/salud_mental/article/view/1257
- López, M. M., Celis, K. F., González-Forteza, C., Saltijeral, M. T., Schiavon, R., Collado, M. E., Ortiz, O., & Lira, L. R. (2019). Relationship between perceived stigma and depressive symptomatology in women who legally interrupt pregnancy in Mexico City. *Salud Mental*, 42, 25-32. <https://doi.org/10.17711/SM.0185-3325.2019.004>
- Major, B., Appelbaum, M., Beckman, L., Dutton, M. A., Russo, N. F., & West, C. (2009). Abortion and mental health: Evaluating the evidence. *American Psychologist*, 64(9), 863-890. <https://doi.org/10.1037/a0017497>
- Maroto-Vargas, A. (2010). Al amparo de la noche: abortos clandestinos y salud mental. *Revista Costarricense de Psicología*, 29(4), 65-75. <http://www.rcps-cr.org/openjournal/index.php/RCPs/article/view/56>
- National Collaborating Centre for Mental Health (2011). *Induced abortion and mental health: A systematic review of the mental health outcomes of induced abortion, including their prevalence and associated factors*. Academy of Medical Royal Colleges. https://www.aomrc.org.uk/wp-content/uploads/2016/05/Induced_Abortion_Mental_Health_1211.pdf
- O'Donnell, A. T., O'Carroll, T., & Toole, N. (2018). Internalized stigma and stigma-related isolation predict women's psychological distress and physical health symptoms post-abortion. *Psychology of Women Quarterly*, 42(2), 220-234. <https://doi.org/10.1177/0361684317748937>
- Oram, S., Khalifeh, H., & Howard, L. M. (2017). Violence against women and mental health. *Lancet Psychiatry*, 4(2), 159-170. [https://doi.org/10.1016/S2215-0366\(16\)30261-9](https://doi.org/10.1016/S2215-0366(16)30261-9)
- Orihuela-Cortés, F., Álvarez del Río, A., Lagunes-Córdoba, R., & Marván, M. L. (2022). Stigma toward women who have had an induced abortion and associated variables. *Women's Reproductive Health*, 1-12. <https://doi.org/10.1080/23293691.2022.2094738>
- Pedersen, W. (2008). Abortion and depression: A population-based longitudinal study of young women. *Scandinavian Journal of Public Health*, 36(4), 424-428. <https://doi.org/10.1177/1403494807088449>
- Radloff, L. S. (1977). The CES-D Scale: A self-report depression scale for research in the general population. *Applied Psychological Measurement*, 1(3), 385-401. <https://doi.org/10.1177/014662167700100306>
- Ramos-Lira L., Celis, K. F., López M. M., Saltijeral M. T., González-Forteza, C., Schiavon R., Collado M. E., & Ortiz, O. (2021). Salud mental e interrupción legal del embarazo: resultados de un estudio exploratorio en la ciudad de México. In Pérez Armendariz E. M., Durante Montiel I., & Figueroa Pérez, M. I. (Eds.), *Ciencia, salud y género* (pp. 156-172). Universidad Nacional Autónoma de México, Facultad de Medicina. <http://libros.facmed.unam.mx/wp-content/uploads/2021/09/Ciencia-salud-y-genero.pdf>
- Ramos-Lira, L., Saltijeral-Méndez, M. T., Romero-Mendoza, M., Caballero-Gutiérrez, M. A., & Martínez-Vélez, N. A. (2001). Violencia sexual y problemas asociados en una muestra de usuarias de un centro de salud. *Salud Pública de México*, 43, 182-91. <https://saludpublica.mx/index.php/spm/article/view/6305>
- Rees, D. I., & Sabia, J. J. (2007). The relationship between abortion and depression: new evidence from the fragile families and child well-being study. *Medical Science Monitor*, 13(10), CR430-436. <https://doi.org/10.12659/msm.502357>
- Roberts, S. C., Biggs, M. A., Chibber, K. S., Gould, H., Rocca, C. H., & Foster, D. G. (2014). Risk of violence from the man involved in the pregnancy after receiving or being denied an abortion. *BMC Medicine*, 12(144). <https://doi.org/10.1186/s12916-014-0144-z>
- Rostagnol, S. (2014). ¿Tolerancia?: debates sobre la legalización del aborto. In Gioscia L. (Ed.), *Más allá de la tolerancia. Ciudadanía y diversidad en el Uruguay contemporáneo* (pp. 170-187). Ediciones Trilce.
- Russo, N. (1976). The motherhood mandate. *Journal of Social Issues*, 32(3), 143-153. <https://doi.org/10.1111/j.1540-4560.1976.tb02603.x>
- Shellenberg, K. M., Moore, A. M., Bankole, A., Juarez, F., Omideyi, A. K., Palomino, N., Sathar, Z., Singh, S., & Tsui, A. O. (2011). Social stigma and disclosure about induced abortion: results from an exploratory study. *Global Public Health*, 6(1), S111-S125. <https://doi.org/10.1080/17441692.2011.594072>
- Singh, S., Remez, L., Sedgh, G., Kwok, L., & Onda, T. (2018). *Abortion worldwide 2017: Uneven progress and unequal access*. Guttmacher. <https://www.guttmacher.org/report/abortion-worldwide-2017>
- Sorhaindo, A. M., Juárez-Ramírez, C., Díaz Olavarrieta, C., Aldaz, E., Mejía Piñeros, M. C., & García, S. (2014). Qualitative evidence on abortion stigma from Mexico City and five states in Mexico. *Women & Health*, 54(7), 622-640. <https://doi.org/10.1080/03630242.2014.919983>
- Steinberg, J. R., & Rubin, L. R. (2014). Psychological aspects of contraception, unintended pregnancy, and abortion. *Policy Insights from the Behavioral and Brain Sciences*, 1(1), 239-247. <https://doi.org/10.1177/2372732214549328>
- Steinberg, J. R., McCulloch, C. E., & Adler, N. E. (2014). Abortion and mental health: Findings from The National Comorbidity Survey-Replication. *Obstetrics & Gynecology*, 123(2), 263-270. <https://doi.org/10.1097/AOG.000000000000092>
- Steinberg, J. R., Tschann, J. M., Furgerson, D., & Harper, C. C. (2016). Psychosocial factors and pre-abortion psychological health: The significance of stigma. *Social Science & Medicine*, 150, 67-75. <https://doi.org/10.1016/j.socscimed.2015.12.007>
- Stotland, N. L. (2019). Update on reproductive rights and women's mental health. *Medical Clinics of North America*, 103(4), 751-766. <https://doi.org/10.1016/j.mcna.2019.02.006>
- Sumner, S. A., Mercy, A. A., Saul, J., Motsa-Nzuza, N., Kwesigabo, G., Buluma, R., Marcelin, L. H., Lina, H., Shawa, M., Moloney-Kitts, M., Kilbane, T., Sommarin, C., Ligiero, D. P., Brookmeyer, K., Chiang, L., Lea, V., Lee, J., Kress, H., Hillis, S. D., & Centers for Disease Control and Prevention (CDC; 2015). Prevalence of sexual violence against children and use of social services - seven countries, 2007-2013. *Morbidity and Mortality Weekly Report*, 64(21), 565-569. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4584766/>
- Turan, J. M., & Budhwani, H. (2021). Restrictive abortion laws exacerbate stigma, resulting in harm to patients and providers. *American Journal of Public Health*, 111(1), 37-39. <https://doi.org/10.2105/AJPH.2020.305998>
- van Ditzhuijzen, J. (2019). Common risk factors for abortion and suicide attempts. *Lancet Psychiatry*, 6(12), 972-973. [https://doi.org/10.1016/S2215-0366\(19\)30443-2](https://doi.org/10.1016/S2215-0366(19)30443-2)
- van Ditzhuijzen, J., ten Have, M., de Graaf, R., Lugtig, P., van Nijntzen, C., & Vollebergh, W. (2017). Incidence and recurrence of common mental disorders after abortion: Results from a prospective cohort study. *Journal of Psychiatric Research*, 84, 200-206. <https://doi.org/10.1016/j.jpsychires.2016.10.006>
- Yin, X., Sun, N., Jiang, N., Xu, X., Gan, Y., Zhang, J., Qiu, L., Yang, C., Shi, X., Chang, J., & Gong, Y. (2021). Prevalence and associated factors of antenatal depression: Systematic reviews and meta-analyses. *Clinical Psychology Review*, 83, 101932. <https://doi.org/10.1016/j.cpr.2020.101932>
- Zia, Y., Mugo, N., Ngure, K., Odoyo, J., Casmir, E., Ayiera, E., Bukusi, E., & Heffron, R. (2021). Psychosocial experiences of adolescent girls and young women subsequent to an abortion in sub-Saharan Africa and globally: A systematic review. *Frontiers Reproductive Health*, 3(638013), 1-13. <https://doi.org/10.3389/frph.2021.638013>